




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07D 243/14, C07C 225/12, 237/06	A1	(11) International Publication Number: WO 00/63186 (43) International Publication Date: 26 October 2000 (26.10.00)
(21) International Application Number: PCT/EP00/03260 (22) International Filing Date: 12 April 2000 (12.04.00) (30) Priority Data: 9908662.1 15 April 1999 (15.04.99) GB (71) Applicant (for all designated States except US): SMITHKLINE BEECHAM PLC [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB). (72) Inventor; and (75) Inventor/Applicant (for US only): HAYES, Jerome, Francis [GB/GB]; SmithKline Beecham Pharmaceuticals, Old Powder Mills, Near Leigh, Tonbridge, Kent TN11 9AN (GB). (74) Agent: VALENTINE, Jill, Barbara; SmithKline Beecham, Corporate Intellectual Property, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: PROCESS AND INTERMEDIATES FOR PREPARING BENZODIAZEPINES <div style="text-align: center;">  (I) </div> (57) Abstract <p>A new process and intermediates for preparing 1,4-benzodiazepines of formula (I) is disclosed: wherein, R¹ is H, OH, Br, I, NR⁵R⁶, CO₂R⁷, CF₃SO₃, C₁₋₄alkoxy, or (4,4'-bipiperidinyl) carbonyl; R² is alkyl, optionally substituted by CO₂R⁷ or CONR⁵R⁶; R³ and R^{3'} are independently H, Ar or C₁₋₆ alkyl optionally substituted by OH, NO₂, NH₂, NR⁵R⁶, halogen, CO₂R⁷, CONR⁵R⁶ or Ar, or together R³ and R^{3'} are =O; R⁴ is H or C₁₋₄ alkyl, optionally substituted by OH, NO₂, NH₂, NR⁵R⁶, halogen, CO₂R⁷, CONR⁵R⁶, or Ar; R⁵ and R⁶ are independently H, Ar, C₁₋₆alkyl, HCO, C₁₋₆alkyl-CO, Ar-CO, C₁₋₆alkyl-SO₂, or Ar-SO₂; and R⁷ is a carboxylic acid protecting group.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakistan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

PROCESS AND INTERMEDIATES FOR PREPARING BENZODIAZEPINES

Field of the Invention

5 This invention relates to processes and intermediates for preparing benzodiazepine vitronectin and fibrinogen antagonists.

Background

10 Tetrahydro-1,4-benzodiazepines form the core structure of a variety of pharmaceutically useful compounds. In particular, WO 93/00095 (PCT/US92/05463), WO 94/14776 (PCT/US93/12436), and WO 95/18619 (PCT/US95/00248) disclose 7-substituted tetrahydro-1,4-benzodiazepines which are reported to be inhibitors of the fibrinogen and vitronectin receptors and are useful as inhibitors of platelet aggregation, osteoporosis, angiogenesis and cancer metastasis.

15 These publications contain a variety of procedures for constructing the basic 1,4-benzodiazepine structure. The prior processes employ expensive reagents and generally require a large number of synthetic steps in the sequence.

 We have now discovered new useful intermediates and a new process for preparing certain 7-substituted 1,4-benzodiazepines. The new process uses a unique


20 cyclization step to form the diazepine ring, and is adaptable to a large scale.

 Teuber et al., Chem. Ber., 98, 2648 (1965), have reported an oxidative method for forming an indole, and Kraus et al., Tet. Lett., 39, 3957 (1998), have reported a cyclization for forming a 3,4-dihydro-1,4-benzodiazepine-2,5-dione, using an acyl hydroquinone and Fremy's salt. Kraus et al., Tet. Lett., 40, 2039


25 (1999), have also reported using Fremy's salt to promote an unusual rearrangement of a 1,4-benzodiazepine-5-one, presumably through a 1-acyl quinone. The compounds of the instant invention do not possess the stabilizing acyl hydroquinone arrangement found in these compounds.

Summary of the Invention

30 In one aspect this invention is a new process for preparing 1,4-benzodiazepines of formula (I):



5 wherein R¹-R⁷ are as defined hereinafter, from substituted 2-hydroxy phenols of formula (II):




10 which comprises reacting a compound of the formula (II) with an oxidizing agent wherein R and R¹-R⁷ are as defined hereinafter.

In another aspect this invention comprises new compounds for preparing 1,4-benzodiazepines.

15 In yet another aspect this process may be used to prepare 2,3,4,5-tetrahydro-7-(4,4'-bipiperidinylcarbonyl)-3-oxo-4-methyl-1H-1,4-benzodiazepine-2-acetic acid, and related active pharmaceutical agents.

Detailed Description of the Invention

20 Many compounds that are useful as pharmaceuticals or intermediates for preparing pharmaceuticals are given by formula (I):



wherein,

25 R¹ is H, OH, Br, I, NR⁵R⁶, CO₂R⁷, CF₃SO₃, C₁-₄alkoxy, or (4,4'-bipiperidinyl)carbonyl;
R² is alkyl, optionally substituted by CO₂R⁷ or CONR⁵R⁶;


R^3 and $R^{3'}$ are independently H, Ar or C_{1-6} alkyl optionally substituted by OH, NO_2 , NH_2 , NR^5R^6 , halogen, CO_2R^7 , $CONR^5R^6$ or Ar, or together R^3 and $R^{3'}$ are =O;

R^4 is H or C_{1-4} alkyl, optionally substituted by OH, NO_2 , NH_2 , NR^5R^6 , halogen, CO_2R^7 , $CONR^5R^6$, or Ar;

R^5 and R^6 are independently H, Ar, C_{1-6} alkyl, HCO, C_{1-6} alkyl-CO, Ar-CO, C_{1-6} alkyl-SO₂, or Ar-SO₂; and

R^7 is a carboxylic acid protecting group.

Compounds of formula (I) may be prepared by a process which comprises,
10 reacting a compound of the formula (II) with an oxidizing agent,



(II)

wherein

R is OH or C_{1-4} alkoxy, and R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^7 are as defined above, and
15 thereafter, treating with a reducing agent.

Preferably R is OH.

Preferably, R^1 is H, OH, NH_2 , bromo, iodo, CO_2R^7 , or C_{1-4} alkoxy. More preferably, R^1 is H or OH.

Preferably, R^2 is $CH_2CO_2R^7$ or $CH_2CONR^5R^6$; more preferably $CH_2CO_2R^7$.

20 Preferably, R^3 and $R^{3'}$ are =O.

Preferably, R^4 is H, C_{1-4} alkyl or Ar- C_{1-4} alkyl. Methyl is especially preferred.

Preferably R^7 is C_{1-4} alkyl, phenyl or benzyl. Methyl, t-butyl or benzyl is especially preferred.


Suitably, R and R^1 are each OH or OCH_3 . Most preferably, R and R^1 are OH.

25 Compounds of formula (II), wherein R, and R^1 - R^4 are as defined above, are useful for the preparation of the compounds of formula (I) and are a feature of this invention. A preferred embodiment comprises compounds of formula (II) wherein R is OH or C_{1-4} alkoxy; R^1 is H, OH, Br, I, C_{1-4} alkoxy, CO_2R^7 , CF_3SO_3 , or (4,4'-bipiperidiny)carbonyl, optionally protected on the basic nitrogen; R^4 is methyl, and
30 R^7 is a carboxy protecting group.

The condensation reaction to form the diazepine ring presumably proceeds through a Schiff's base-type intermediate, as illustrated in Scheme 1, so the oxidant used must be

Scheme 1


5



sufficient to promote oxidation of the aromatic hydroxyl group to a carbonyl and induce the free amino group to condense with it. Salts of ferricyanide and nitrosodisulfonate are useful, especially potassium ferricyanide and potassium nitrosodisulfonate. Agents of similar oxidation potential are also useful, such as silver (I) oxide, thallium (IV) nitrate, cerium (IV) ammonium nitrate, hypervalent iodine compounds, palladium on charcoal with a hydrogen acceptor (e.g., nitrobenzene), acid dichromate, lead tetra-acetate, and potassium periodate. Potassium nitrosodisulfonate (Fremy's salt) is a preferred reactant. Acetic acid and hydrochloric acid are preferred acids for use in the reaction, although other organic acids and mineral acids may also be suitable.

It will be apparent that when R¹ is a hydroxyl group, the reaction may proceed through a quinone-type intermediate, as indicated by the compounds of formula (III), wherein R², R³ and R⁴ are as defined in formula (I).

20




For instance, a preferred compound is (3*S*)-methyl 3-amino-4-(*N*-methyl-*N*-(1,4-benzoquinonylmethyl)carbamoyl) butanoate,


Reduction of the double bond completes the construction of the 2,3,4,5 tetrahydro-1,4-benzodiazepine ring system. Reduction may be accomplished readily
 5 by catalytic hydrogenation, such as by hydrogenation over palladium or platinum, although other means, such as transfer hydrogenation, for instance using ammonium formate and Pd/C in refluxing methanol, may be suitable.

In cases where further unsaturation is present in the ring system, tautomerisation of the Schiff's base may occur. For instance, when there is a 3-oxo
 10 group and a 2-acetic acid group, the double bond may tautomerize, as indicated in Scheme 2.

Scheme 2




Thus, 1,4-benzodiazepine ring systems that have 3-oxo, 2-acetic acid
 15 substituents tautomerize to a 2-ethyldiene configuration, and compounds of formula (IV):



(IV)


wherein R¹ is H, OH, Br, I, NR⁵R⁶, CO₂R⁷, CF₃SO₃, C₁-₄alkoxy, or (4,4'-bipiperidiny)carbonyl; R⁴ is H or C₁-₄ alkyl, optionally substituted by OH, NO₂, NH₂, NR⁵R⁶, halogen, CO₂R⁷, CONR⁵R⁶, or Ar; R⁵ and R⁶ are independently H,
 25 Ar, C₁-₆alkyl, HCO, C₁-₆alkyl-CO, Ar-CO, C₁-₆alkyl-SO₂, or Ar-SO₂; and R⁷ is a carboxylic acid protecting group, are useful intermediates for preparing pharmaceutically active compounds, such as those disclosed in WO 95/18619 (PCT/US95/00248); WO 93/00095 (PCT/US/92/05463) and WO 94/14776 (PCT/US93/12436). Preferably R¹ is H or OH, and R⁴ is methyl.

In a preferred embodiment, this invention is a process for preparing the active pharmaceutical product 2,3,4,5-tetrahydro-7-(4,4'-bipiperidinylicarbonyl)-3-oxo-4-methyl-1H-1,4-benzodiazepine-2-acetic acid, comprising reacting a compound of the formula (V):



wherein R¹ is H, OH, I, Br, CO₂R⁷, CF₃SO₂, C₁-₄alkoxy, or (4,4'-bipiperidinylicarbonyl, optionally protected on one nitrogen; R⁴ is methyl, and R⁷ is a carboxy protecting group; with a reducing agent; and, if necessary, converting R¹ to (4,4'-bipiperidinylicarbonyl, and, if necessary, removing any protecting groups. It will be apparent that the α,β unsaturated ester may be reduced at any point in the synthesis subsequent to the cyclization reaction. Reduction may be accomplished, for instance, using palladium or platinum on carbon. Preferably, R¹ is H or OH.

In another preferred embodiment, this invention is a process for preparing 2,3,4,5-tetrahydro-7-(4,4'-bipiperidinylicarbonyl)-3-oxo-4-methyl-1H-1,4-benzodiazepine-2-acetic acid, comprising treating the compound of formula (VI):




wherein R¹ is H, OH, I, Br, CO₂R⁷, CF₃SO₃, C₁-₄alkoxy, or (4,4'-bipiperidinylicarbonyl, optionally protected on the basic nitrogen; R⁴ is methyl; and R⁷ is a carboxy protecting group; with an oxidizing agent, and, in any order, treating with a reducing agent and, if necessary, converting R¹ to (4,4'-bipiperidinylicarbonyl; and, if necessary, removing any protecting groups. Preferably, R¹ is H or OH, and the product is converted to 2,3,4,5-tetrahydro-7-(4,4'-bipiperidinylicarbonyl)-3-oxo-4-methyl-1H-1,4-benzodiazepine-2-acetic acid via a palladium catalyzed aminocarbonylation.

It will be understood that one may use a variety of functional groups in the process for R¹, and subsequently transform such groups to the (4,4'-bipiperidinyl)carbonyl sidechain. For instance, one may convert a hydroxyl group at R¹ to a triflate group, and add the bipiperidinylcarbonyl group via a palladium catalyzed aminocarbonylation reaction, as described in WO 97/24336 (PCT/IB96/01502). It will of course be evident that an alkoxy substituent may be converted to a hydroxy substituent, such as by hydrolysis, and further transformed by similar means to the desired 4,4'-bipiperidinylcarbonyl group. One may also use an aminocarbonylation reaction to convert the bromo or iodo group to the bipiperidinylcarbonyl amino group. Additionally, a hydrogen at R¹ may be converted to bromo or iodo group, as described for instance in WO 97/24336 (PCT/IB96/01502), and converted via an aminocarbonylation reaction to the bipiperidinylcarbonyl group. An ester group may be hydrolyzed to a carboxyl group and reacted with the bipiperidinyl group using conventional amide bond forming reagents, such as dicyclohexylcarbodiimide or via an acid chloride.

As is apparent from the foregoing, compounds according to formula (VI), are useful for preparing compounds of formula (V), and constitute a part of this invention.

Scheme 3 illustrates some of the representative methods for preparing 2,3,4,5-tetrahydro-7-(4,4'-bipiperidinylcarbonyl)-3-oxo-4-methyl-1H-1,4-benzodiazepine-2-acetic acid.

Scheme 3



- (a) MeNH₂ (8.03M solution in EtOH), toluene (b) 10% Pd-C, EtOH, H₂ (c) N-Cbz-aspartic acid β-methyl ester, DCC, optionally HOBt; (d) ammonium cerium(IV) nitrate, MeCN, H₂O; (e) H₂, 10% Pd-C, ethanol; (f) potassium nitrosodisulfonate chloroform, acetic acid, HCl; (g) H₂, 10% Pd-C, ethanol; (h) potassium nitrosodisulfonate chloroform, acetic acid, HCl; (i) H₂, 10% Pd-C, ethanol, H₃PO₄

The meaning of any substituent at any one occurrence in formulas (I)-(VI) or any subformula thereof is independent of its meaning, or any other substituent's meaning, at any other occurrence, unless specified otherwise.

- Abbreviations and symbols commonly used in the chemical arts are used herein to describe the compounds, reactions and reagents of this invention.

Certain radical groups are abbreviated herein. t-Bu refers to the tertiary butyl radical, Boc refers to the t-butyloxycarbonyl radical, Ph refers to the phenyl radical, CBZ or Z refers to the benzyloxycarbonyl radical, Bn refers to the benzyl radical, Me refers to methyl, Et refers to ethyl, and Ac refers to acetyl.

- Certain reagents are abbreviated herein. DCC refers to dicyclohexylcarbodiimide, DMAP refers to dimethylaminopyridine, DIEA refers to diisopropylethylamine. THF refers to tetrahydrofuran, DMF refers to dimethylformamide, Pd-C refers to a palladium on carbon catalyst, TEA refers to triethylamine, TFA refers to trifluoroacetic acid.

- C₁₋₄alkyl as applied herein is meant to include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl and t-butyl. C₁₋₆alkyl additionally includes pentyl, n-pentyl, isopentyl, neopentyl and hexyl and the simple aliphatic isomers thereof. Any C₁₋₄alkyl or C₁₋₆alkyl group may be optionally substituted by halo, -OR', SR', -CN, -NR'R', -NO₂, -CF₃, CF₃S(O)_r-, -CO₂R', -CONR'₂, C₃₋₆cycloalkyl, or Ar, unless otherwise indicated, where R' is H or C₁₋₄alkyl. A substituent on a C₁₋₆ alkyl may be on any carbon atom which results in a stable structure, and is available by conventional synthetic techniques.

- Ar, or aryl, as applied herein, means phenyl or naphthyl, or phenyl or naphthyl substituted by one to three moieties. In particular, such moieties may be C₁₋₄alkyl, C₁₋₄alkoxy, C₁₋₄alkylthio, trifluoromethyl, OH, F, Cl, Br or I.

- The reactive functional groups of each synthetic fragment are suitably protected as known in the art. Suitable protective groups are disclosed in Greene, PROTECTIVE GROUPS IN ORGANIC CHEMISTRY, John Wiley and Sons, New York, 1981. For example, the Boc, Cbz, phthaloyl or Fmoc group may be used for protection of an amino (or the nitrogen of a piperidiny). The Boc or Cbz group is

generally preferred for protection of an amino group, such as on the basic nitrogen of a 4,4'-bipiperidinyl group. The term the basic nitrogen of the 4,4'-bipiperidinyl group, as used herein, indicates the nitrogen which is not attached to the benzodiazepine nucleus via a carbonyl group.

5 A methyl, ethyl, t-Bu, cHex, benzyl, substituted benzyl, (pivaloyl)methyl or (2-methyl-2-methoxypropanoyl)methyl ester may be used for the protection of the carboxyl group. A suitably substituted carbobenzyloxy group or benzyl group may be also be used for the hydroxyl group or amino group. Suitable substitution of the carbobenzyloxy or benzyl protecting groups is ortho and/or para substitution with
10 chloro, bromo, nitro, methoxy or methyl, and is used to modify the reactivity of the protective group. C₁₋₆alkoxy, phenyl and benzyl are conventional carboxy protecting groups.

 Methods for removal of the a carboxy or amino protecting group are well known in the art. For example, an alkyl or cycloalkyl ester may be removed by
15 basic hydrolysis, for instance an alkali metal hydroxide, such as sodium, potassium or lithium hydroxide in a suitable solvent, such as aqueous alcohol. A benzyl ester is typically removed by hydrogenation over a palladium catalyst. A basic nitrogen protected by a t-butyloxycarbonyl group, or a t-butyl ester, is typically removed by acid treatment, such as by trifluoroacetic acid or hydrochloric acid, optionally
20 diluted with a solvent, such as methylene chloride and/or dioxane. The benzyloxycarbonyl group is generally removed by hydrogenation over a palladium catalyst. A trifluoroacetyl group is typically removed by basic hydrolysis, such as by treatment with an alkali metal hydroxide in a suitable solvent. One useful synthetic method for protecting the basic nitrogen of a piperidinyl group is to carry
25 the group through a synthesis as a pyridinyl group, which may be reduced with a platinum or palladium catalyst toward the end of the synthesis to remove the protecting group.

 The simple starting materials for preparing the compounds of this invention are commercially available or prepared by routine methods well known in the art.

30 The intermediate compounds of this invention are useful as intermediates in the preparation of pharmaceutically active compounds, in particular compounds which have fibrinogen and vitronectin antagonist properties.

General

Nuclear magnetic resonance spectra were recorded at 270 MHz. CDCl₃ is deuteriochloroform, DMSO-d₆ is hexadeuteriodimethylsulfoxide, and CD₃OD is tetradeuteriomethanol. Chemical shifts are reported in parts per million (δ) downfield from the internal standard tetramethylsilane. Abbreviations for NMR data are as follows: s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet, dd = doublet of doublets, dt = doublet of triplets, app = apparent, br = broad. J indicates the NMR coupling constant measured in Hertz. HRMS indicates high resolution mass spectroscopy.

EXAMPLES

Example 1

15 Preparation of 2,3,4,5-tetrahydro-7-(N-CBZ-4,4'-bipiperidinylcarbonyl)-3-oxo-4-methyl-2-methoxycarbonylethylidenyl)-1H-1,4-benzodiazepine (9)

a) N-(2,5-dimethoxybenzylidene) methylamine.

To a solution of 2,5-dimethoxybenzaldehyde (20.00g, 0.12 mol) in toluene (100 ml) was added methylamine (22.5 ml of an 8.03M solution in ethanol, 0.18 mol) at room temperature. The solution became cloudy within 10 min. due to the condensation of water. The reaction had gone to completion within 30 min. according to GC analysis and the solvent was then removed in vacuo to give the required product; (21.17g, 98%) as a yellow oil. δH (CDCl₃, 270 MHz) 3.50 (3H, s), 3.80 (3H, s), 3.85 (3H, s), 6.80 (1H, m), 7.00 (1H, m), 7.50 (1H, m) and 8.70 (1H, s).

b) methyl (2,5-dimethoxybenzyl) amine (1)

A solution of N-(2,5-dimethoxybenzylidene) methylamine (21.17g, 0.12 mol) in ethanol (200 ml) was hydrogenated over 10% palladium on charcoal (water content 50%, 4.0g). The reaction had gone to completion within 2h according to GC analysis. The catalyst was filtered off and the filtrate was concentrated to dryness. The title compound was thus obtained as a yellow oil (18.00g, 85%); δH (CDCl₃, 270 MHz) 1.70 (1H, s br), 2.45 (3H, s), 3.70 (2H, s), 3.75 (3H, s) 3.80 (3H, s), 6.70 - 6.85 (3H, m).

c) (3*S*)-methyl 3-benzyloxycarbonylamino-4-(*N*-methyl-*N*-2,5-dimethoxybenzylcarbamoyl) butanoate (2)

To a solution of methyl (2,5-dimethoxybenzyl) amine (10.00g, 55.2mmol) and *N*-CBZ aspartic acid β -methyl ester (17.10g, 60.9mmol) in ethyl acetate (100 ml) was added dicyclohexylcarbo diimide (12.50g, 60.7mmol) at room temperature. A thick precipitate formed within 15 min. and after 1h the reaction mixture was filtered. The filtrate was concentrated to dryness to afford the crude required product as a thick oil and as a mixture of rotamers (30.73g, 100%) δ H (CDCl₃, 270 MHz) 2.60 - 2.85 (2H, m), 2.90 (1.5H, s), 3.10 (1.5H, s), 3.65 (2H, s), 3.70 (3H, s), 3.80 (3H, s), 4.50 - 4.75 (1H, m), 5.05 (1H, d), 5.10 (1H, s), 4.70 (0.5H, d), 5.80 (0.5H, d), 6.70 - 6.80 (3H, m) and 7.25 - 7.40 (5H, m).

d) (3*S*)-methyl 3-benzyloxycarbonylamino-4-(*N*-methyl-*N*-(1,4-benzoquinonylmethyl) carbamoyl) butanoate (3)

To a solution of (3*S*)-methyl 3-benzyloxycarbonylamino-4-(*N*-methyl-*N*-2,5-dimethoxybenzylcarbamoyl) butanoate (3) (47.0g, 0.106mol) in acetonitrile (2.35L) was added cerium IV ammonium nitrate (145.5g, 0.265mol) in water (454 ml) at -14°C -10°C. The reaction mixture became dark brown-orange immediately and the colour became lighter over 1h. The temperature was allowed to rise to 5 - 10°C over 2h. and then to 12 - 15°C over 90 min. The solvent was then removed in vacuo and the residue was dissolved in dichloromethane (500 ml) and was washed with water. The aqueous layer was back extracted with dichloromethane (500 ml). The combined organic extracts were dried (Mg SO₄) and concentrated to dryness. The residue was chromatographed on silica gel using ethyl acetate as eluant. Hence the title compound was obtained as a viscous yellow oil (44.5g, 100%); δ H (CDCl₃, 270 MHz) 2.60 - 3.00 (2H, m), 3.25 (3H, s), 3.70 (3H, s), 3.85 (1H, m), 4.40 (2H, ABq), 5.10 (2H, m), 5.60 (1H, d br), 6.55 (1H, m), 6.75 (2H, m) and 7.40 (5H, m).

e) (3*S*)-methyl 3-amino-4-(*N*-methyl-*N*-(2,5-dihydroxybenzyl) carbamoyl) butanoate (4)

A solution of (3*S*)-methyl 3-benzyloxycarbonylamino-4-(*N*-methyl-*N*-(1,4-benzoquinonylmethyl) carbamoyl)butanoate (44.5g, 0.100mol) in ethanol (1.50L) was hydrogenated over 10% palladium on charcoal (16.0g) at room temperature under one atmosphere of hydrogen for 16h. The catalyst was filtered off after this time and the filtrate was concentrated to dryness. (The crude product was not characterised but was used directly in the next step). δ H (d-6 DMSO, 270 MHz

40°C) 2.40 (1H, dd), 2.70 (1H, dd), 2.90 (1.5H, s), 3.05 (1.5H, s), 3.30 (2H, s br), 3.60 (3H, s), 4.35 (1H, ABq), 4.50 (1H, ABq), 6.45 (2H, m) and 6.60 (1H, m).

5 f) 2,3,4,5-tetrahydro-7-hydroxy-3-oxo-4-methyl-2-(methoxycarbonylethylidenyl)-1H-1,4-benzodiazepine (5)

To a solution of (3*S*)-methyl 3-amino-4-(*N*-methyl-*N*-(2,5-dihydroxybenzyl) carbamoyl) butanoate (4) (30.30g, 0.107mmol) in water (2.08L) and acetic acid (233 ml) and chloroform (5.0L) was added hydrochloric acid (9.7 ml of an 11.0M aqueous solution, 0.107mmol) followed by Fremy's salt (63.4g, 0.237mol) in water (3.0L) at room temperature. The reaction mixture was stirred for 3h after which time the aqueous layer was separated. The chloroform layer was washed with water (1.0L) and was then concentrated to dryness. The resultant dark foamy residue was chromatographed on silica gel using ethyl acetate:hexane - 3:1 as eluant. Hence the title compound was obtained as white solid (7.29g, 26%). δ H (CDCl₃, 270 MHz) 3.00 (3H, s), 3.75 (3H, s), 4.25 (2H, s br), 5.40 (1H, s), 5.80 (1H, s), 6.70 - 6.80 2H, m), 6.85 (1H, d) and 10.50 (1H, s). High Resolution Mass Spectrum (HRMS): Found M^+H 263.1016. C₁₃H₁₅N₂O₄ requires 263.1032.

20 g) 2,3,4,5-tetrahydro-7-(trifluoromethanesulfonyloxy-3-oxo-4-methyl-2-methoxycarbonylethylidenyl)-1H-1,4-benzodiazepine (6)

To a solution of 2,3,4,5-tetrahydro-7-hydroxy-3-oxo-4-methyl-2-(methoxycarbonylethylidenyl)-1H-1,4-benzodiazepine (5) (1.00g, 3.82mmol) in pyridine (3.0 ml) was added trifluoromethanesulfonic anhydride (7.06 μ l, 4.21mmol) at 0°C. The reaction mixture was stirred for 90 min. after which time it was diluted with ether (100 ml) and washed with water (250 ml). The aqueous layer was extracted with further ether (ca 100 ml) and the combined organic layers were dried (MgSO₄) and concentrated to dryness. The residue was chromatographed on silica gel using ethyl acetate:hexane - 2:3 as eluant. Hence the title compound was obtained as a white solid (1.10g, 73%) m/z 395 [M+H]⁺, 363 [M - MeOH+H]⁺, 335 [M - HCO₂Me+H]⁺; δ H (CDCl₃, 400 MHz) 3.10 (3H, s), 3.75 (3H, s), 4.30 (2H, s br), 5.50 (1H, s), 7.10 (1H, d), 7.15 (1H, d), 7.20 (1H, dd) and 10.70 (1H, s); HRMS: Found 394.0431. C₁₄H₁₃F₃N₂O₆S requires 394.0446.

h) 2,3,4,5-tetrahydro-7-(*N*-CBZ-4,4'-bipiperidinylcarbonyl)-3-oxo-4-methyl-2-methoxycarbonylethylidenyl)-1*H*-1,4-benzodiazepine (7)

A 50 ml 3-necked flask containing a mixture of 2,3,4,5-tetrahydro-7-(trifluoromethanesulfonyloxy-3-oxo-4-methyl-2-methoxycarbonylethylidenyl)-1*H*-1,4-benzodiazepine (6) (0.27g, 0.68mmol), diisopropylethylamine (1.4 ml, 8.05mmol), *N*-CBZ-4,4'-bipiperidine hydrochloride (298 mg, 0.88mmol), palladium II chloride bistrisphenylphosphine (10 mg, 0.014mmol) and water (30μl, 1.66mmol) in *N*-methylpyrrolidinone (2.2 ml) was evacuated and filled with carbon monoxide gas several times. The reaction mixture was then heated to 90 - 95°C under an atmosphere of carbon monoxide with vigorous stirring. After 5h palladium (0) tetrakis(trisphenylphosphine) (100mg) was added and the carbonylation was continued. The reaction had gone to completion according to HPLC analysis after 16h. A solution assay of the reaction mixture against a reference standard indicated the yield to be 181mg, 46%. The reaction mixture was diluted with dichloromethane (30 ml) and was filtered. The filtrate was washed with water (2 x 30 ml) and was then concentrated to dryness in vacuo. The residue was chromatographed on silica gel using ethyl acetate as eluant to give the title compound as a pale yellow foam (144mg, 37%). This was triturated with ether (ca 4 ml) to give a purified sample as a white powder. m/z 575 $[M+H]^+$; δH ($CDCl_3$, 400 MHz) 1.10 - 1.45 (6H, m), 1.60 - 1.85 (4H, m), 2.75 (4H, m), 3.15 (3H, s), 3.75 (3H, s), 3.85 (1H, s v br), 4.15 - 4.30 (4H, m), 4.70 (1H, s br), 5.15 (2H, s), 5.45 (1H, s), 7.00 (1H, d), 7.30 - 7.40 (7H, m) and 10.70 (1H, s).

i) methyl 2,3,4,5-tetrahydro-7-(4,4'-bipiperidinylcarbonyl)-3-oxo-4-methyl-1*H*-1,4-benzodiazepine-2-acetate (8)

A suspension of 2,3,4,5-tetrahydro-7-(*N*-CBZ-4,4'-bipiperidinylcarbonyl)-3-oxo-4-methyl-2-methoxycarbonylethylidenyl)-1*H*-1,4-benzodiazepine (7) (375 mg, 0.653mmol) in methanol (20 ml) and water (10 ml) was acidified to pH 2.5 with a 26% aqueous solution of phosphoric acid. The resultant mixture was hydrogenated over 10% palladium on charcoal (Johnson Matthey type 87L, 375 mg) at 60 psi in a Parr Shaker at room temperature. The reaction had gone to completion within 2h according to HPLC analysis. The catalyst was filtered off and the filtrate was concentrated in vacuo to ca 5 ml. Water (10 ml) was added and the pH was then adjusted to 10 with a 35% aqueous solution of ammonia. A white precipitate formed and the suspension was cooled to 5°C for 16h. The precipitate was then filtered off

and dried in the open atmosphere to give the title compound as a white powder (257mg, 89%).

j) 2,3,4,5-tetrahydro-7-(4,4'-bipiperidinylcarbonyl)-3-oxo-4-methyl-1H-1,4-benzodiazepine-2-acetic acid (9)

The compound of Example 1(i) (0.26 mmol) was dissolved in methanol (9 ml), and 1.0 N sodium hydroxide (0.81 ml, 0.81 mmol) was added. The solution was stirred overnight at room temperature, and concentrated. The residue was dissolved in water/acetonitrile (3 ml), cooled to 0°C, acidified with HCl, and concentrated to yield the title compound.

Example 2

Preparation of 2,3,4,5-tetrahydro-7-(4,4'-bipiperidinylcarbonyl)-3-oxo-4-methyl-1H-1,4-benzodiazepine-2-acetic acid (12)

a) N-(2,5-hydroxybenzylidene) methylamine

Using the procedure of Example 1(a), except substituting 2,5-dihydroxybenzaldehyde for 2,5-dimethoxybenzaldehyde, the title compound was prepared. δ H (CDCl₃, 270 MHz), 3.50 (3H,s), 6.75 (1H,m), 6.85 (2H,m) and 8.25 (1H,s)

b) N-methyl, N-(2,5-dihydroxybenzyl) amine (21)

Using the procedure of Example 1(b), except substituting the compound of Example 2(a) for the compound of Example 1(a), the title compound was prepared. δ H (CDCl₃, 270 MHz) 2.25 (3H,s), 3.60 (2H,s) and 6.50 (3H,m).

c) (3S)-methyl 3-benzyloxycarbonylamino-4-(N-methyl-N-2,5-dihydroxybenzylcarbamoyl) butanoate (22)

Using the procedure of Example 1(c), except substituting the compound of Example 2(b), and chromatographing the crude product over silica gel using ethyl acetate:hexane – 1:1 – 1:0 as eluent, the title compound was prepared (46%). δ H (CDCl₃, 270 MHz) 2.65 (1H,dd), 2.80 (1H,dd), 2.95 (3H,s), 3.60 (3H,s), 4.35 (2H, ABQ), 5.05 (2H,m), 5.80 (1H,d), 6.65 (1H,s), 6.75 (2H,m), 7.25 - 7.50 (5H,m) and 8.60 (2H,s(br)).

e) (3*S*)-methyl 3-amino-4-(*N*-methyl-*N*-(2,5-dihydroxybenzyl) carbamoyl) butanoate (23)

A solution of 22 (7.82g, 18.80 mmol) in absolute ethanol 107ml was hydrogenated over 10% palladium on charcoal (0.78g) for 20h. The catalyst was filtered off and the filtrate was concentrated to dryness. The residue was chromatographed on silica gel using ethyl acetate : dichloromethane : methanol - 4:4:2 as eluant. The title compound was obtained as a colourless foam (3.6g, 65%).

f) 2,3,4,5-tetrahydro-7-hydroxy-3-oxo-4-methyl-2-(methoxycarbonyl ethylidenyl)-1*H*-1,4-benzodiazepine (24)

Using the procedure of Example 1(f), except substituting the compound of Example 2(e) for the compound of Example 1(e), the title compound was prepared.

g) 2,3,4,5-tetrahydro-7-(4,4'-bipiperidinylcarbonyl)-3-oxo-4-methyl-1*H*-1,4-benzodiazepine-2-acetic acid (9)

Using the compound of Example 2(f) and following the procedures of Examples 1(d) - 1(j), the title compound is prepared.

Example 3

20 Preparation of 2,3,4,5-tetrahydro-7-(4,4'-bipiperidinylcarbonyl)-3-oxo-4-methyl-1*H*-1,4-benzodiazepine-2-acetic acid

a) *N*-(2-hydroxybenzylidene) methylamine

Using the procedure of Example 1(a), except substituting 2-hydroxy-benzaldehyde for 2,5-dimethoxy-benzaldehyde, the title compound was prepared. δ H (CDCl₃, 270MHz) 6.80 (1*H*,t), 7.00 (1*H*,s), 7.20-7.30 (2*H*,m) and 8.40 (1*H*,s).

b) *N*-methyl, *N*-(2-hydroxybenzyl) amine (31)

Using the procedure of Example 1(b), except substituting the compound of Example 3(a) for the compound of Example 1(a), the title compound was prepared. δ H (CDCl₃, 270MHz) 2.50 (3*H*,s), 4.00 (2*H*,s) 6.75 (1*H*,t), 6.80 (1*H*,d), 7.00 (1*H*,d) and 7.20 (1*H*,t).

c) (3*S*)-methyl 3-benzyloxycarbonylamino-4-(*N*-methyl-2-hydroxybenzylcarbamoyl) butanoate (32)

Using the procedure of Example 1(c), except substituting the compound of Example 3(b) for the compound of Example 1(b) and adding HOBT (1.1 equiv), and chromatographing the crude product using ethyl acetate:hexane – 4:6 – 1:0 as eluent, the title compound was prepared (82%). δ H (CDCl₃, 270MHz) 2.65 (1H,dd), 2.80 (1H,dd), 3.20 (3H,s), 3.60 (3H,s), 4.40 (2H,ABQ), 5.10 (2H,m), 5.70 (1H,d), 6.80 (1H,t), 6.90 (1H,d), 7.15 (1H,dd), 7.25 (1H,dt) and 9.05 (1H,s).

10 d) (3*S*)-methyl 3-amino-4-(*N*-methyl-*N*-(2-hydroxybenzyl) carbamoyl) butanoate (33)

A solution of 32 (11.26g 28.15mmol) in absolute ethanol (78ml) was hydrogenated over 3% palladium on charcoal (2.62g) at room temperature for 2h. The catalyst was filtered off and the filtrate was concentrated to dryness. The residue was chromatographed on silica gel using ethyl acetate: chloroform: methanol – 9:9:2 as eluant. The title compound was obtained as a colourless foam (6.90g, 92%). δ H (CDCl₃, 270MHz) 2.55 (1H,dd), 2.70 (1H,dd), 3.20 (3H,s), 3.65 (3H,s), 4.15 (1H,t), 4.50 (2H,ABQ), 6.80 (1H,t), 6.95 (1H,d), 7.15 (1H,d) and 7.25 (1H,t). [M + H]⁺ 267

20 Using the procedure of Example 1(f), except substituting the compound of Example 3(e), the title compound is prepared.

f) methyl 2,3,4,5-tetrahydro-3-oxo-4-methyl-1*H*-1,4-benzodiazepine-2-acetate

Following the procedure of Example 4, except substituting the compound of Example 3(e), the title compound is prepared.

g) methyl 2,3,4,5-tetrahydro-7-(4,4'-bipiperidinylcarbonyl)-3-oxo-4-methyl-1*H*-1,4-benzodiazepine-2-acetate (9)

Following the procedures set forth in WO 97/24336 (PCT/IB96/01502), the title compound is prepared.

Example 4

Preparation of methyl 2,3,4,5-tetrahydro-7-hydroxy-3-oxo-4-methyl-1*H*-1,4-benzodiazepine-2-acetate (10)

35 A solution of the compound of Example 1(f) (1.73g, 6.6mmol) in absolute ethanol (173ml) was hydrogenated over 10% palladium on charcoal (1.73g) for 16h.


at room temperature. The catalyst was then filtered off and the filtrate was concentrated to dryness. The residue was slurried in dichloromethane; ether - 1:1 and the product was then filtered off as a white solid (800mg, 46%). δ H (DMSO, 270MHz) 2.60(1h, dd), 2.80(1H, dd), 2.90 (3H, s), 3.60(3H,s), 3.75(1H, d),
5 4.80(1H,m), 5.10(1H,d), 5.35(1H, d), 6.40(3H, m) and 8.55(1H, s).

The above description fully discloses how to make and use the present invention. However, the present invention is not limited to the particular
10 embodiments described hereinabove, but includes all modifications thereof within the scope of the following claims. The various references to journals, patents and other publications which are cited herein comprises the state of the art and are incorporated herein by reference as though fully set forth.

15

Claims:

1. A process for preparing a compound of the formula (I):



(I)

wherein,

R^1 is H, OH, Br, I, NR^5R^6 , CO_2R^7 , CF_3SO_3 , C_{1-4} alkoxy, or (4,4'-bipiperidiny)carbonyl;

R^2 is alkyl, optionally substituted by CO_2R^7 or $CONR^5R^6$;

R^3 and $R^{3'}$ are independently H, Ar or C_{1-6} alkyl optionally substituted by OH, NO_2 , NH_2 , NR^5R^6 , halogen, CO_2R^7 , $CONR^5R^6$ or Ar, or together R^3 and $R^{3'}$ are =O;


R^4 is H or C_{1-4} alkyl, optionally substituted by OH, NO_2 , NH_2 , NR^5R^6 , halogen, CO_2R^7 , $CONR^5R^6$, or Ar;

R^5 and R^6 are independently H, Ar, C_{1-6} alkyl, HCO, C_{1-6} alkyl-CO, Ar-CO, C_{1-6} alkyl- SO_2 , or Ar- SO_2 ; and

R^7 is a carboxylic acid protecting group,

which comprises

reacting a compound of the formula (II) with an oxidizing agent,



(II)

wherein

R is OH or C_{1-4} alkoxy, and R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^7 are as defined above, and,

thereafter, treating with a reducing agent.

2. A process according to claim 1 wherein R^2 is $CH_2CO_2R^7$, and together R^3 and $R^{3'}$ are =O.


3. A process according to claim 1 wherein R^1 is H, OH or C_{1-4} alkoxy.

4. A process according to claim 1 wherein the oxidizing agent is potassium ferricyanide or potassium nitrosodisulfonate.

5

5. A process according to claim 1 wherein the reducing agent is hydrogen in the presence of a palladium or platinum on carbon catalyst.

6. A compound of formula (III):



10

(III)

wherein,

R^2 is alkyl, optionally substituted by CO_2R^7 or $CONR^5R^6$;

15 R^3 and $R^{3'}$ are independently H, Ar or C_{1-6} alkyl optionally substituted by OH, NO_2 , NH_2 , NR^5R^6 , halogen, CO_2R^7 , $CONR^5R^6$ or Ar, or together R^3 and $R^{3'}$ are =O;

R^4 is H or C_{1-4} alkyl, optionally substituted by OH, NO_2 , NH_2 , NR^5R^6 , halogen, CO_2R^7 , $CONR^5R^6$, or Ar;


20 R^5 and R^6 are independently H, Ar, C_{1-6} alkyl, HCO, C_{1-6} alkyl-CO, Ar-CO, C_{1-6} alkyl-SO₂, or Ar-SO₂; and

R^7 is a carboxylic acid protecting group,

7. A compound according to claim 6 which is (3*S*)-methyl 3-amino-4-(*N*-methyl-*N*-(1,4-benzoquinonylmethyl) carbamoyl) butanoate.

25

8. A method for preparing 2,3,4,5-tetrahydro-7-(4,4'-bipiperidinylcarbonyl)-3-oxo-4-methyl-1*H*-1,4-benzodiazepine-2-acetic acid comprising, in any order, reacting a compound of formula (V):



(V)

wherein

R^1 is H, OH, Br, I, C_{1-4} alkoxy, CO_2R^7 , CF_3SO_2 , or (4,4'-bipiperidiny)carbonyl,

5 optionally protected on one nitrogen;


R^4 is methyl, and

R^7 is a carboxy protecting group;

with a reducing agent, and, if necessary, converting R^1 to (4,4'-bipiperidiny)carbonyl; and

10 if necessary, removing any protecting groups.

9. A compound of the formula:



(V)

15 wherein

R^1 is H, OH, I, Br, C_{1-4} alkoxy, CO_2R^7 , CF_3SO_2 , or (4,4'-bipiperidiny)carbonyl optionally protected on the basic nitrogen;

R^4 is methyl, and

R^7 is a carboxy protecting group.

20

10. A compound according to claim 8, which is:

2,3,4,5-tetrahydro-3-oxo-4-methyl-2-(methoxycarbonyl ethylidenyl)-1H-1,4-benzodiazepine;


2,3,4,5-tetrahydro-7-hydroxy-3-oxo-4-methyl-2-(methoxycarbonylethylidenyl)-1H-1,4-benzodiazepine;

25

2,3,4,5-tetrahydro-7-(trifluoromethanesulfonyloxy-3-oxo-4-methyl-2-methoxycarbonylethylidenyl)-1H-1,4-benzodiazepine; or

2,3,4,5-tetrahydro-7-(*N*-CBZ-4,4'-bipiperidiny)carbonyl)-3-oxo-4-methyl-2-methoxycarbonylethylidenyl)-1H-1,4-benzodiazepine.

11. A method for preparing 2,3,4,5-tetrahydro-7-(4,4'-bipiperidinylcarbonyl)-3-oxo-4-methyl-1H-1,4-benzodiazepine-2-acetic acid, comprising reacting a compound of the formula:




5

(VI)

wherein R¹ is H, OH, Br, I, C₁₋₄alkoxy, CO₂R⁷, CF₃SO₃, or (4,4'-bipiperidinyl)carbonyl, optionally protected on one nitrogen; R⁴ is methyl, and

10 R⁷ is a carboxy protecting group;
with an oxidizing agent, and, in any order, treating with a reducing agent; and, if necessary, converting R¹ to (4,4'-bipiperidinyl)carbonyl; and removing any protecting groups.

15 12. A compound of the formula:



(II)

wherein

20 R is OH or C₁₋₄alkoxy;
R¹ is H, OH, Br, I, C₁₋₄alkoxy, CO₂R⁷, CF₃SO₃, or (4,4'-bipiperidinyl)carbonyl, optionally protected on the basic nitrogen;;
R⁴ is methyl, and
R⁷ is a carboxy protecting group.

25 13. A compound according to claim 11, which is:
(3*S*)-methyl 3-amino-4-(*N*-methyl-*N*-(2,5-dihydroxybenzyl) carbamoyl) butanoate;
or
(3*S*)-methyl 3-amino-4-(*N*-methyl-*N*-(2-hydroxybenzyl) carbamoyl) butanoate.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 00/03260

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D243/14 C07C225/12 C07C237/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	J. F. HAYES: "An Alternative Synthesis of a Potent GPIIb/IIIa Receptor Antagonist" SYNLETT, no. S1, 1999, pages 865-6, XP000915438 schemes 2, 3; pages 865, right-hand column to page 866, left-hand column ----- -/-	1-3, 5, 6, 8, 9, 11, 12

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

14 July 2000

Date of mailing of the international search report

26/07/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Hass, C

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 00/03260

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	MILLER W H ET AL: "Enantiospecific Synthesis of SB 214857, a Potent, Orally Active, Nonpeptide Fibrinogen Receptor Antagonist" TETRAHEDRON LETTERS,NL,ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, vol. 36, no. 52, 25 December 1995 (1995-12-25), pages 9433-9436, XP004026650 ISSN: 0040-4039 page 9433, compounds 3, 4; page 9434, scheme 1; page 9435, scheme 2 ----	1,2,6,9, 11,12
A	J. M. SAMANEN ET AL.: "Potent, Selective, Orally Active 3-Oxo-1,4-benzodiazepine GPIIb/IIIa Integrin Antagonists" JOURNAL OF MEDICINAL CHEMISTRY, vol. 39, no. 25, 6 December 1996 (1996-12-06), pages 4867-70, XP002142556 page 4869, scheme 1 ----	1,2,6,9, 11,12
A	WO 93 00095 A (SMITHKLINE BEECHAM CORP.) 7 January 1993 (1993-01-07) cited in the application page 21, scheme 2 ----	1,6,9,12
A	WO 94 14776 A (SMITHKLINE BEECHAM CORP.) 7 July 1994 (1994-07-07) cited in the application page 14, scheme 2 ----	1,6,9,12
A	R. M. KEENAN ET AL.: "Conformational Preferences in a Benzodiazepine Series of Potent Nonpeptide Fibrinogen Receptor Antagonists" JOURNAL OF MEDICINAL CHEMISTRY, vol. 42, no. 4, 25 February 1999 (1999-02-25), pages 545-59, XP002142557 tables 1,2 ----	9
A	WO 95 18619 A (SMITHKLINE BEECHAM CORP.) 13 July 1995 (1995-07-13) cited in the application abstract ----	9
A	WO 95 17888 A (SMITHKLINE BEECHAM CORP.) 6 July 1995 (1995-07-06) page 12, compound 5 claim 1 -----	12

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 00/03260

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9300095 A	07-01-1993	AU 666318 B	08-02-1996
		AU 2271192 A	25-01-1993
		EP 0593603 A	27-04-1994
		JP 6509074 T	13-10-1994
		NZ 243326 A	26-10-1995
		PT 100631 A,B	30-09-1993
		US 5693636 A	02-12-1997
		ZA 9204760 A	31-03-1993
WO 9414776 A	07-07-1994	AU 689762 B	09-04-1998
		AU 5873694 A	19-07-1994
		CA 2152631 A	07-07-1994
		CN 1094718 A,B	09-11-1994
		CN 1215054 A	28-04-1999
		EP 0674623 A	04-10-1995
		JP 8505145 T	04-06-1996
		NZ 259802 A	24-10-1997
		NZ 314922 A	25-11-1998
		SG 63579 A	30-03-1999
		ZA 9309515 A	24-08-1995
WO 9518619 A	13-07-1995	AP 542 A	01-10-1996
		AU 699186 B	26-11-1998
		AU 1600195 A	01-08-1995
		BG 100698 A	28-02-1997
		BG 102863 A	31-05-1999
		BR 9506499 A	09-09-1997
		CN 1143323 A	19-02-1997
		CZ 9601981 A	11-06-1997
		EP 0738150 A	23-10-1996
		HU 76282 A	28-07-1997
		JP 9508620 T	02-09-1997
		NO 962859 A	05-09-1996
		NZ 279394 A	27-05-1998
		SG 64884 A	25-05-1999
		SK 86996 A	05-02-1997
		ZA 9500045 A	23-01-1996
WO 9517888 A	06-07-1995	AU 1441195 A	17-07-1995
		EP 0738144 A	23-10-1996
		JP 9507846 T	12-08-1997
		ZA 9410318 A	01-08-1995